

Conservation laws for self-adjoint first order evolution equation

Igor Leite Freire

Centro de Matemática, Computação e Cognição
Universidade Federal do ABC - UFABC
Rua Catequese, 242, Jardim, 09090 – 400
Santo André, SP - Brasil
 E-mail: igor.freire@ufabc.edu.br

Abstract

In this work we consider the problem on group classification and conservation laws of the general first order evolution equations. We obtain the subclasses of these general equations which are quasi-self-adjoint and self-adjoint. By using the recent new conservation theorem due to Ibragimov, we establish the conservation laws of the equations admitting self-adjoint equations. We illustrate our results applying them to the inviscid Burgers' equation. In particular an infinite number of new symmetries of these equations are found and their corresponding conservation laws are established.

2000 AMS Mathematics Classification numbers:

76M60, 58J70, 70G65

Key words: Lie point symmetry, Ibragimov's Theorem, conservation laws, inviscid Burgers equation.

1 Introduction

The Lie point symmetries of evolution equations with one spatial variable

$$u_t = F\left(t, x, u, \frac{\partial u}{\partial x}, \dots, \frac{\partial^n u}{\partial x^n}\right), \quad (1)$$

with $n \geq 2$, has been studied by many authors, see [2, 5, 6, 11, 12, 13, 14, 16, 18, 25, 26, 31, 32] and references therein.

For example, if $n = 2$, equation (1) includes the nonlinear heat equation, the Burgers equation, Fokker-Planck equation, Black-Scholes equation and, more generally, reaction-diffusion-convection equations, see [6, 12, 16, 25, 31, 32].

The Korteweg-de Vries (KdV) equation, cylindrical KdV and modified KdV are examples of third order evolution equations, see [2, 16].

When $n = 4$, equation (1) includes the modified Kuramoto-Sivashinsky equation, the Cahn-Hilliard equation, the thin film equation and others, see [5, 13, 26].

However, the first order equation

$$u_t + f(t, x, u, u_x) = 0 \quad (2)$$

seems to have received few attention.

To the best of our knowledge, the earliest work involving first order evolution equations and Lie point symmetries was [24], where the authors studied equation (2) with $f = a(u)u_x$, for some particular functions $a(u)$. After it, Nadjafikhah [19] obtained projectable symmetries of equation

$$u_t + a(u)u_x = 0 \quad (3)$$

and in [20] the same author classifies the similarity solutions of the symmetries obtained in [19]. Equation (3) is known as inviscid Burgers equation.

The purpose of this work is to deal with the problem on group classification of the general first order evolution equation and how to obtain conservation laws from the Lie point symmetries. We intend to

- find the first order evolution equations that admit (quasi) self-adjoint equations;
- obtain close formulae to express conservation laws for equations of the type (2) using recent results due to Ibragimov [16];
- generalize the previous results on group classification of equation (3);
- establish conservation laws for equation (3).

The paper is organized as the follows. In the next section we obtain the general determining equations for the components of symmetry generators, the (quasi) self-adjointness condition and establish the corresponding conservation laws for self-adjoint equations of the type (2). We also obtain new Lie point symmetry generators of (3) and some conservation laws for it are established.

2 Main results

In this section we shall consider the group classification problem and how to find conservation laws for equation (2) with $f_{u_x} \neq 0$. Hereafter all functions will be assumed to be smooth, the summation over the repeated indices is understood, $u_x = \frac{\partial u}{\partial x}$ and $u_t = \frac{\partial u}{\partial t}$.

Following the standard Lie approach [1, 15, 23], let

$$X = \tau(t, x, u) \frac{\partial}{\partial t} + \xi(t, x, u) \frac{\partial}{\partial x} + \eta(t, x, u) \frac{\partial}{\partial u} \quad (4)$$

be a Lie point symmetry generator of equation (2). Then the coefficients τ , ξ and η satisfy the following equation

$$\begin{aligned} & \eta_t - \xi_t u_x + (\tau_t - \eta_u + \xi_u u_x) f + \xi f_x + \tau f_t + \eta f_u - \tau_u f^2 \\ & + (\eta_x + \eta_u u_x - \xi_x u_x - \xi_u u_x^2) f_{u_x} + (\tau_x + \tau_u u_x) f f_{u_x} = 0. \end{aligned} \quad (5)$$

We observe that (5) is one equation to be solved for 4 unknown functions ξ , τ , η and f .

To obtain the full symmetry group of equation (2) it is necessary to find all possible functions ξ , τ , η , f satisfying the relation (5). Thus, a complete group classification of (2) is impossible.

Let us now consider the problem of finding conservation laws for equations of the type (2).

If an equation possesses variational structure, the Noether Theorem can be employed in order to establish conservation laws for it, *e.g.*, see [3, 4, 10, 22].

However, it is well-known that evolution equations do not possess variational structure. Then, they cannot be obtained from the Euler-Lagrange equations and the Noether's Theorem cannot be applied to them in order to obtain conservation laws.

Fortunately there are some alternative methods to obtain conservation laws for equations without Lagrangians: the direct method, the characteristic method, the variational approach, the symmetry conditions, the direct construction formula and the partial Noether approach. For a more detailed discussion, see [2, 16, 22]. Although some of these methods could be employed in order to establish conservation laws for equation (3), in this paper we shall use recent results due to Ibragimov [16] in order to construct conservation laws for equations of the type (2). We shall refer to the new conservation theorem established in [16] (see Theorem 3.5 in the reference) as Ibragimov's Theorem.

The Ibragimov's Theorem on conservation laws can be summarized by the following algorithm (see [16] for more details): given a PDE

$$F = F(x, u, \partial u, \dots, \partial^n u) = 0,$$

where $\partial^k u$ denotes the set of all k th order derivatives of u ,

- we construct a Lagrangian $\mathcal{L} = vF$.
- From the Euler-Lagrange equations, the following system is obtained:

$$F(x, u, \partial u, \dots, \partial^n u) = 0, \quad (6)$$

$$F^*(x, u, v, \partial u, \partial v, \dots, \partial^n u, \partial^n v) = 0. \quad (7)$$

The second equation of the system (6) - (7) is called adjoint equation to $F = 0$.

Equation (7) is said to be quasi-self-adjoint if the system (6) - (7) is equivalent to the original equation (6) upon the substitution $v = \varphi(u)$ such that $\varphi'(u) \neq 0$, *i.e.*,

$$F^*(x, u, v, \partial u, \partial v, \dots, \partial^n u, \partial^n v)|_{v=\varphi(u)} = \phi F(x, u, \partial u, \dots, \partial^n u), \quad (8)$$

for some function $\phi = \phi(x, u, \partial u, \dots, \partial^n u)$.

If equation (8) is true with $\varphi(u) = u$, then (6) is said to be self-adjoint. For more details, see [5, 16, 17].

- The conserved vector is $C = (C^i)$, where

$$\begin{aligned} C^i &= \xi^i \mathcal{L} + W \left[\frac{\partial \mathcal{L}}{\partial u_i} - D_j \left(\frac{\partial \mathcal{L}}{\partial u_{ij}} \right) + D_j D_k \left(\frac{\partial \mathcal{L}}{\partial u_{ijk}} \right) - \dots \right] \\ &\quad + D_j(W) \left[\frac{\partial \mathcal{L}}{\partial u_{ij}} - D_k \left(\frac{\partial \mathcal{L}}{\partial u_{ijk}} \right) + \dots \right] + \dots \end{aligned} \quad (9)$$

and $W = \eta - \xi^j u_j$.

Then, applying this algorithm to equation (2), we obtain:

- Lagrangian:

$$\mathcal{L} = vu_t + vf(t, x, u, u_x). \quad (10)$$

- Adjoint equation: the adjoint equation to (2) is $F^* = 0$, where

$$F^* = -v_t - v_x f_{u_x} + v f_u - v f_{xu_x} - v u_x f_{uu_x} - v f_{u_x u_x} u_{xx}. \quad (11)$$

- Components of the conserved vector $C = (C^0, C^1)$:

$$\begin{aligned} C^0 &= (\eta + \tau f - \xi u_x) v, \\ C^1 &= (\eta + \tau f - \xi u_x) v f_{u_x}. \end{aligned} \quad (12)$$

2.1 Quasi-self-adjointness condition of equation (2)

Supposing that

$$F^*|_{v=\varphi(u)} = \phi F,$$

where $F = u_t + f(t, x, u, u_x)$ and F^* is given by (11), we obtain $\phi = -\varphi'(u)$ and

$$\begin{cases} f_{u_x u_x} = 0, \\ \varphi(u) f_u - \varphi'(u) f_{u_x} u_x - \varphi(u) f_{uu_x} u_x - \varphi(u) f_{xu_x} + \varphi'(u) f = 0. \end{cases} \quad (13)$$

From (13) we conclude that $f = \alpha(t, x, u) u_x + \beta(t, x, u)$, with $\alpha \neq 0$. Hence, the functions $\varphi(u)$, $\alpha(t, x, u)$ and $\beta(t, x, u)$ should satisfy

$$\beta_u \varphi(u) + \varphi'(u) \beta = \varphi(u) \alpha_x. \quad (14)$$

It follows that if (2) is quasi-self-adjoint, we have two cases to consider:

Case 1: If $\beta \neq 0$, in order for (14) to be true, then

$$\frac{\alpha_x - \beta_u}{\beta} = \frac{\varphi'(u)}{\varphi(u)}, \quad (15)$$

and in this case

$$\varphi(u) = c \exp \int \frac{\alpha_x - \beta_u}{\beta} du, \quad (16)$$

where $c \in \mathbb{R}$ is an arbitrary constant.

Case 2: If $\beta = 0$, from (14), $\alpha = \alpha(t, u)$ and φ is an arbitrary function.

Reciprocally, if $f = \alpha(t, u)u_x$ in (2), it is easy to check that (2) is quasi-self-adjoint. When $f = \alpha(t, x, u)u_x + \beta(t, x, u)$, then f satisfies (13) if (15) is satisfied and then, function φ is given by (16).

According to equation (12), taking $v = \varphi(u)$ (quasi-self-adjointness condition), a conservation law for equation

$$u_t + \alpha(t, x, u)u_x + \beta(t, x, u) = 0, \quad (17)$$

is $D_t C^0 + D_x C^1 = 0$, where α and β are as considered in cases 1 or 2, and

$$\begin{aligned} C^0 &= [\eta + \tau\beta + (\tau\alpha - \xi)u_x] \varphi(u), \\ C^1 &= [\eta\alpha + \xi\beta - (\tau\alpha - \xi)u_t] \varphi(u). \end{aligned} \quad (18)$$

2.2 Self-adjointness condition of equation (2)

Let us now find the class of the self-adjoint equations of the type (2). Whenever $\varphi = u$, equation (14) becomes

$$\beta_u u + \beta = u \alpha_x.$$

Again we have two cases to consider:

Case 1: If $\beta \neq 0$, then

$$\beta = \frac{1}{u} \int u \alpha_x du + \frac{\lambda(t, x)}{u}, \quad (19)$$

for some function $\lambda = \lambda(t, x)$.

Case 2: If $\beta = 0$ then $\alpha = \alpha(t, u)$.

From equation (19), we observe that the case $\beta = 0$ occurs if and only if $\alpha_x = \lambda = 0$. Then, the most general form of a self-adjoint equation of the type (2) is (17), where β is given by (19).

It is easy to check that all equations of the type (17), with β satisfying (19), are self-adjoints.

Equation (17) includes

- inviscid Burgers equation, taking $\alpha = a(u)$ and $\beta = 0$, see [24, 19, 20, 7];
- linear transport equation, taking $\alpha = q(x)$ and $\beta = 0$, see [8, 9].

From equation (18), taking $\varphi = u$ (self-adjointness condition), a conservation law for equation (17) is $D_t C^0 + D_x C^1 = 0$, where

$$\begin{aligned} C^0 &= [\eta + \tau\beta + (\tau\alpha - \xi)u_x] u, \\ C^1 &= [\eta\alpha + \xi\beta - (\tau\alpha - \xi)u_t] u. \end{aligned} \quad (20)$$

2.3 Theorems on self-adjoint equations and conservation laws

Our main results on the self-adjointness conditions and conservation laws can be summarized by the following theorems.

Theorem 1. *Let (4) be a Lie point symmetry generator of equation (2). Then the symmetry coefficients satisfy (5).*

Corollary 1. *The determining equations of (17) are given by*

$$\begin{aligned} \eta_t + \beta(\tau_t - \eta_u) + \beta_x \xi + \beta_t \tau + \beta_u \eta - \beta^2 \tau_u + \alpha \eta_x + \alpha \beta \tau_x &= 0, \\ -\xi_t + \alpha \tau_t + \beta \xi_u + \alpha_x \xi + \alpha_t \tau + \alpha_u \eta - \alpha \beta \tau_u - \alpha \xi_x + \alpha^2 \tau_x &= 0. \end{aligned} \tag{21}$$

Proof Substituting $f = \alpha(t, x, u)u_x + \beta(t, x, u)$ into (5), we obtain (21). $\square$

Theorem 2. *The following statements about equation (17) are true:*

1. *If $\beta = 0$, (17) is quasi-self-adjoint if and only if $\alpha = \alpha(t, u)$.*
2. *If $\beta \neq 0$, (17) is quasi-self-adjoint if and only if the functions α and β satisfy the relation (15), for some function $h = h(u)$, and φ is given by (16).*

Theorem 3. *Equation (2) is self-adjoint if and only if $f = \alpha(t, u)u_x$ or $f = \alpha u_x + \beta$, where β is given by (19).*

Theorem 4. *A conservation law for the system*

$$\begin{cases} u_t + f(t, x, u, u_x) = 0, \\ -v_t - v_x f_{u_x} + v f_u - v f_{xu_x} - v u_x f_{uu_x} - v f_{u_x u_x} u_{xx} = 0. \end{cases}$$

is $\text{Div}(C) = D_t C^0 + D_x C^1 = 0$, where C^0 and C^1 are given by (12) and τ, ξ and η are the coefficients of the generator (4).

Theorem 5. *A conservation law for equation (17), with α and β as in Theorem 3, is $\text{Div}(C) = D_t C^0 + D_x C^1 = 0$, where C^0 and C^1 are given by (20) and τ, ξ and η are the coefficients of the generator (4).*

2.4 Inviscid Burgers equation

We recall that to obtain the group classification of (2) we need to construct all possible functions ξ, τ, η and f obeying (5). As mentioned above, this is an underdetermined problem and a general group classification is impossible.

Regarding equation (17), from Corollary 1, we conclude that it possesses an infinity dimensional symmetry Lie algebra.

Equation (17) covers the so-called inviscid Burgers equation and it describes turbulence phenomena, for instance, compressible gas dynamics, shallow water flow, weather prediction, plasma modelling, rarefied gas dynamics and many others, see [7, 8, 9, 19, 20, 21, 27, 28, 29, 30].

In [7] the random Riemann problem for Burgers equation is solved. In [8] a numerical scheme to approximate the m th moment of the solution of the one-dimensional random linear transport equation

is studied. In [9] a numerical scheme for the random linear transport equation is presented. In [30] numerical methods are employed for solving hyperbolic conservation laws.

Distributional products and solutions of the inviscid Burgers equation

$$u_t + uu_x = 0 \quad (22)$$

are studied in [27, 28]. In [27] the concept of global α -solution for equation (22) is introduced, as well as the existence of “delta-soliton” travelling waves. In [28] new solutions are presented. System of transport equations are considered in [21, 29]. Further details can be found in the references cited above. In what follows, the Lie point symmetries and conservation laws for (3) shall be discussed.

2.4.1 Projectable symmetries of inviscid Burgers equation

Let us now consider the symmetries of equation (3).

From equation (5), the symmetry coefficients τ , ξ and η satisfy the following determining equations

$$\eta_t + a(u)\eta_x = 0, \quad (23)$$

$$\eta a'(u) + \tau_x a(u)^2 - \xi_t + \tau_t a(u) - \xi_x a(u) = 0. \quad (24)$$

Since the system (23) - (24) is underdetermined, we use the symmetries obtained by Nadjafikhah in [19]. The *ansatz* employed by Nadjafikhah in [19] was to consider the projectable symmetries of (3). For more details, see [19].

Supposing that $\tau = \tau(t, x)$ and $\xi = \xi(t, x)$, Nadjafikhah obtained the following basis to the symmetry Lie algebra (see [19, 20]):

$$\begin{aligned} X_1 &= \frac{\partial}{\partial t}, \quad X_2 = \frac{\partial}{\partial x}, \quad X_3 = x \frac{\partial}{\partial x} + t \frac{\partial}{\partial t}, \quad X_4 = t \frac{\partial}{\partial t} - \frac{a(u)}{a'(u)} \frac{\partial}{\partial u}, \\ X_5 &= t \frac{\partial}{\partial x} + \frac{1}{a'(u)} \frac{\partial}{\partial u}, \quad X_6 = x \frac{\partial}{\partial t} - \frac{a(u)^2}{a'(u)} \frac{\partial}{\partial u}, \\ X_7 &= t^2 \frac{\partial}{\partial t} + tx \frac{\partial}{\partial x} + \frac{x - ta(u)}{a'(u)} \frac{\partial}{\partial u}, \\ X_8 &= tx \frac{\partial}{\partial t} + x^2 \frac{\partial}{\partial x} + \frac{a(u)(x - ta(u))}{a'(u)} \frac{\partial}{\partial u}. \end{aligned} \quad (25)$$

Two questions naturally arise:

Q1: Are there symmetries such that $(\xi_u, \tau_u) \neq (0, 0)$?

Q2: Could it possible to find symmetries more general than that obtained in [19]?

An (simple) answer to **Q1** is the following: suppose $\tau = \tau(u)$ and $\xi = \xi(u)$. From (23) - (24) we conclude that $\eta = 0$ and

$$X = \tau(u) \frac{\partial}{\partial t} + \xi(u) \frac{\partial}{\partial x}$$

is a Lie point symmetry generator of (3).

In the next subsection **Q2** shall be answered.

2.4.2 New Lie point symmetry generators for equation (3)

With regard to **Q2**, according to Corollary 1, if $\beta = 0$ in (17) and

$$X = \tau(t, x) \frac{\partial}{\partial t} + \xi(t, x) \frac{\partial}{\partial x} + \eta(t, x, u) \frac{\partial}{\partial u} \quad (26)$$

is a projectable symmetry generator of (17), then the determining equations (5) do not have terms involving derivatives of the Lie point symmetry generators with respect to u . Thus it is easy to check that the components of the vector field

$$X_\lambda = \lambda(u) \tau(t, x) \frac{\partial}{\partial t} + \lambda(u) \xi(t, x) \frac{\partial}{\partial x} + \lambda(u) \eta(t, x, u) \frac{\partial}{\partial u},$$

where $\lambda = \lambda(u)$ is a smooth function, satisfy the determining equations (21). So, the field X_λ is a nonprojectable Lie point symmetry generator and the following results are immediate consequences of the Corollary 1, Theorem 3 and Theorem 5.

Theorem 6. *Let X be a projectable Lie point symmetry generator of equation*

$$u_t + \alpha(t, u) u_x = 0, \quad (27)$$

and $\lambda(u)$ a smooth function. Then the vector field $X_\lambda = \lambda(u)X$ is a Lie point symmetry of the equation (27).

Corollary 2. *Let (26) be a projectable Lie point symmetry generator of equation (27), $\lambda(u)$ a smooth function and*

$$\begin{aligned} C^0 &= \lambda(u) [\eta + (\tau\alpha - \xi) u_x] u, \\ C^1 &= \lambda(u) [\eta - (\tau\alpha - \xi) u_t] u. \end{aligned} \quad (28)$$

Then the vector field $C = (C^0, C^1)$ is a conserved field for equation (27).

Remarks:

1. According to Theorem 6, given a projectable symmetry generator X of equation (27), from it we can construct an infinite number of nonprojectable symmetry generators, given by $X_\lambda = \lambda(u)X$, where $\lambda = \lambda(u)$ is a smooth function.

2. From Corollary 2 it is easy to conclude that given a projectable symmetry generator X of equation (27), it is possible to obtain an infinite number of conservation laws, given by formulae (28).

2.5 Conservation laws for inviscid Burgers equation

Here we shall use the Ibragimov's Theorem on conservation laws [16] to establish the conservation laws for equation (3).

From Theorem 3, equation (3) is self-adjoint and the Theorem 5 can be employed in order to establish conservation laws for it.

From (28) and (3) a conserved vector is $C = (C^0, C^1)$, where

$$\begin{aligned} C^0 &= [\eta + (\tau a(u) - \xi) u_x] u, \\ C^1 &= [\eta a(u) - (\tau a(u) - \xi) u_t] u. \end{aligned} \quad (29)$$

With regard to the time and spatial translational invariance, it is easy to check that the conservation laws are trivial. Let $A(u)$ be a function such that

$$A'(u) = u a(u). \quad (30)$$

For the symmetry X_3 , the conservation law is $D_t C^0 + D_x C^1 = 0$, where:

$$C^0 = [ta(u) - x] u u_x,$$

$$C^1 = [x - ta(u)] u u_t.$$

Equation (3) is a first-order equation. Consequently, zero-order conservation laws are more important than first-order one. So we intend to simplify the components C^0 and C^1 in order to establish zero-order conservation laws for (3).

Since $C^0 = D_x(tA) - xD_x(u^2/2)$ and $C^1 = D_t(xu^2/2) - tD_t(A)$, the conserved vector $C = (C^0, C^1)$ can be simplified using the fact

$$\begin{aligned} D_t C^0 + D_x C^1 &= D_x(A) + tD_t D_x(A) - xD_t D_x\left(\frac{u^2}{2}\right) \\ &\quad + D_t\left(\frac{u^2}{2}\right) + xD_x D_t\left(\frac{u^2}{2}\right) - tD_x D_t(A) \\ &= D_t\left(\frac{u^2}{2}\right) + D_x(A). \end{aligned}$$

It follows that

$$C^0 = \frac{u^2}{2}, \quad C^1 = A, \quad (31)$$

where A is given in (30), provides a conserved vector for equation (3).

Below we present, in a schematic form, the conservation laws associated to the Lie point symmetry generators $X_4, \dots, X_8$. First we present the conservation laws given by Theorem 5. In the following, we give the simplified vector employing the same procedure used in order to obtain the components (31).

1. For the symmetry X_4 , the components of the vector field given by Theorem 5 are

$$C^0 = -\frac{a(u)}{a'(u)} u + t a(u) u u_x, \quad C^1 = -\frac{a(u)^2}{a'(u)} u - t a(u) u u_t.$$

The simplified components are

$$C^0 = \frac{a u}{a'}, \quad C^1 = \frac{a^2 u}{a'} - A. \quad (32)$$

2. For the symmetry X_5 , the components of the vector field given by Theorem 5 are

$$C^0 = \frac{u}{a'(u)} - t u u_x, \quad C^1 = \frac{a(u)}{a'(u)} u + t u u_t.$$

The simplified components are

$$C^0 = \frac{u}{a'}, \quad C^1 = \frac{a u}{a'} - \frac{u^2}{2}. \quad (33)$$

3. For the symmetry X_6 , the components of the vector field given by Theorem 5 are

$$C^0 = -\frac{a(u)^2}{a'(u)}u + x a(u) u u_x, \quad C^1 = -\frac{a(u)^3}{a'(u)}u - x a(u) u u_t.$$

The simplified components are

$$C^0 = \frac{a(u)^2}{a'(u)}u + A, \quad C^1 = \frac{a(u)^3}{a'(u)}u. \quad (34)$$

4. For the symmetry X_7 , the components of the vector field given by Theorem 5 are

$$C^0 = \frac{x - t a(u)}{a'(u)}u + (t^2 a(u) - t x) u u_x, \quad C^1 = \frac{x - t a(u)}{a'(u)}a(u)u - (t^2 a(u) - t x) u u_t.$$

The simplified components are

$$C^0 = \frac{x - t a(u)}{a'(u)}u + \frac{t u^2}{2}, \quad C^1 = \frac{x - t a(u)}{a'(u)}a(u)u + 2 t A - \frac{x u^2}{2}. \quad (35)$$

5. For the symmetry X_8 , the components of the vector field given by Theorem 5 are

$$C^0 = \frac{x - t a(u)}{a'(u)}a(u)u + (t x a(u) - x^2) u u_x, \quad C^1 = \frac{x - t a(u)}{a'(u)}a(u)^2 u - (t x a(u) - x^2) u u_t.$$

The simplified components are

$$C^0 = \frac{x - t a(u)}{a'(u)}a(u)u + x u^2 - t A, \quad C^1 = \frac{x - t a(u)}{a'(u)}a^2(u)u + x A. \quad (36)$$

3 Conclusion

In this paper we considered the general problem on group classification of the general first order evolution equation (2). We found the general classes of the quasi-self and self-adjoint equations of the type (2). By using the recent Ibragimov's Theorem on conservation laws, we derive the general formulae to the conserved fields. Our main results are summarized in theorems 1 - 6, corollaries 1, 2 and in the conservation laws for inviscid Burgers equation established in section 3 (equations (31) - (36)).

We believe that the research on group analysis of equations type (17) can be promising. From Theorem 3, this equation is the most general first order evolution equation that admits self-adjoint equations. Following the Ibragimov's Theorem on conservation laws [16], we have a closed form to express its conservation laws given by equation (29).

From (17) and the determining equations (21) it is noted that we can obtain an underdetermined system of equations to be solved for 5 unknown functions α , β , τ , ξ and η . Thus, the set of symmetries is infinity.

With regard to the inviscid Burgers equation (3), the set of the determining equations is also underdetermined, as we can observe in (23)-(24). See also [19, 24]. From Theorem 6 an infinite

number of new symmetries of equation (3) are presented supposing that $X_\lambda = \lambda(u)X$, where $\lambda(u)$ is a smooth function, and

$$X = \tau(t, x) \frac{\partial}{\partial t} + \xi(t, x) \frac{\partial}{\partial x} + \eta(t, x, u) \frac{\partial}{\partial u}$$

is a projectable symmetry generator of (3). This just is one more possible ansatz to determine Lie point symmetries of (3). From Corollary 2, the corresponding conservation laws associated to the symmetries given by Theorem 6 are established.

A natural question that arises is: which more ansatz can someone use in order to obtain more general symmetries of the inviscid Burgers equation? This is a question that, hopefully, can inspire some more progress in this area.

Acknowledgments

I would like to thank the referee of this paper. The referee not only read this article very carefully, but also contributed with numerous comments/corrections that improved the paper.

I am grateful to Priscila L. da Silva for her careful reading of this paper.

References

- [1] G. W. Bluman and S. Kumei, *Symmetries and Differential Equations* (Springer, New York, 1989).
- [2] G. Bluman, Temuerchaolu and S. C. Anco, New conservation laws obtained directly from symmetry action on a known conservation law, *J. Math. Anal. Appl.* **322**(1) (2006) 233–250.
- [3] Y. D. Bozhkov and I. L. Freire, Conservations laws for critical Kohn-Laplace equations on the Heisenberg group, *J. Nonlinear Math. Phys.*, **15**(1) (2008) 35–47.
- [4] Y. Bozhkov and I. L. Freire, Special conformal groups of a riemannian manifold and the Lie point symmetries of the nonlinear Poisson equations, *J. Diff. Equ.*, **249**(4) (2010) 872–913.
- [5] M. S. Bruzón, M. L. Gandarias and N. H. Ibragimov, Self-adjoint sub-classes of generalized thin film equations, *J. Math. Anal. Appl.* **357**(1) (2009) 307–313.
- [6] R. Cherniha, M. Serov and I. Rassokha, Lie symmetries and form-preserving transformations of reaction-diffusion-convection equations, *J. Math. Anal. Appl.*, **342**(2) (2008) 1363–1379.
- [7] M. C. C. Cunha and F. A. Dorini, Statistical moments of the solution of the random Burgers Riemann problem, *Math. Comput. Simulat.*, **79**(5) (2009) 1440–1451.
- [8] F. A. Dorini and M. C. C. Cunha, Statistical moments of the random linear transport equation, *J. Comput. Phys.* **227**(19) (2008) 8541–8550.
- [9] F. A. Dorini and M. C. C. Cunha, A finite volume method for the mean of the solution of the random transport equation, *Appl. Math. Comput.*, **187**(2) (2007) 912–921.
- [10] I. L. Freire, On the paper “Symmetry analysis of wave equation on sphere” by H. Azad and M. T. Mustafa, *J. Math. Anal. Appl.*, **367**(2) (2010) 716–720.

- [11] I. L. Freire, Self-adjoint sub-classes of third and fourth-order evolution equations, submitted.
- [12] M. L. Gandarias, M. Torrisi and R. Tracinà, On some differential invariants for a family of diffusion equations, *J. Phys. A: Math. Theor.* **40**(30) (2007) 8803–8813.
- [13] M. L. Gandarias and N. H. Ibragimov, Equivalence group of a fourth-order evolution equation unifying various non-linear models, *Commun. Nonlinear Sci. Numer. Simul.* **13**(2) (2008) 259–268.
- [14] R. Gazizov and N. Ibragimov, Lie symmetry analysis of differential equations in finance, *Nonl. Dyn.* **17**(4) (1998) 387–407.
- [15] N. H. Ibragimov, *Transformation groups applied to mathematical physics*, (D. Reidel Publishing Co., Dordrecht, 1985).
- [16] N. H. Ibragimov, A new conservation theorem, *J. Math. Anal. Appl.* **333**(1) (2007) 311–328.
- [17] N. H. Ibragimov, Quasi-self-adjoint differential equations, *Archives of ALGA* (2007).
- [18] V. I. Lahno and A. M. Samoilenko, Group classification of nonlinear evolution equations. I. Invariance under semisimple local transformation groups, *Differ. Equ.* **38**(3) (2002) 384–391.
- [19] M. Nadjafikhah, Lie symmetries of inviscid Burgers equation, *Adv. Appl. Clifford Alg.* **19**(1) (2009) 101–112.
- [20] M. Nadjafikhah, Classification of similarity solutions for inviscid Burgers equation, *Adv. Appl. Clifford Alg.*, **20**(1) (2009) 71–77.
- [21] M. Nedeljkov and M. Oberguggenberger, Interactions of delta shock waves in a strictly hyperbolic system of conservation laws, *J. Math. Anal. Appl.* **344**(2) (2008) 1143–1157.
- [22] R. Naz, F. M. Mahomed and D. P. Mason, Comparison of different approaches to conservation laws for some partial differential equations in fluid mechanics, *Appl. Math. Comput.* **205**(1) (2008) 212–230.
- [23] P. J. Olver, *Applications of Lie groups to differential equations* (Springer, New York, 1986).
- [24] A. Ouhadan and E. H. El Kinani, Lie symmetries of the equation $u_t(x, t) + g(u)u_x(x, t) = 0$, *Adv. Appl. Clifford Alg.* **17**(1) (2007) 95–106.
- [25] R. O. Popovych and N. M. Ivanova, New results on group classification of nonlinear diffusion-convection equations, *J. Phys. A: Math. Gen.* **37**(30) (2004) 7547–7565.
- [26] C. Qu, Symmetries and solutions to the thin film equations, *J. Math. Anal. Appl.* **317**(2) (2006) 381–397.
- [27] C. O. R. Sarrico, Distributional products and global solutions for nonconservative inviscid Burgers equation, *J. Math. Anal. Appl.* **281**(2) (2003) 641–656.
- [28] C. O. R. Sarrico, New solutions for the one-dimensional nonconservative inviscid Burgers equation, *J. Math. Anal. Appl.* **317**(2) (2006) 496–509.

- [29] C. Shen, M. Sun, Interactions of delta shock waves for the transport equations with split delta functions, *J. Math. Anal. Appl.* **351**(2) (2009) 747–755.
- [30] Y. H. Zahran, Central ADER schemes for hyperbolic conservation laws, *J. Math. Anal. Appl.* **346**(1) (2008) 120–140.
- [31] R. Zhdanov and V. Lahno, Group classification of the general evolution equation: local and quasilocal symmetries, *SIGMA* **1** (2005).
- [32] R. Zhdanov and V. Lahno, Group classification of the general second-order evolution equation: semi-simple invariance groups, *J. Phys. A: Math. Theor.* **40**(19) (2007) 5083–5103.